

garp.org.ng



GARP



GARP INTERNATIONAL JOURNAL OF HEALTH SCIENCES



garp.org.ng



Vol. 1, Issue 3, Pp. 13-24; AUG., 2026

ASSESSMENT OF QUALITY OF MATERNAL AND NEWBORN CARE AND ITS DETERMINANTS IN PRIMARY HEALTH CARE FACILITIES IN NIGERIA: IMPLICATIONS FOR SERVICE DELIVERY IMPROVEMENT

¹NWANEKEZI, Adaobi Fidelia & ²ASIBOR, Raphael Ehikhuemhen PhD

¹College of postgraduate Studies, Faculty of Applied health sciences, Department of Nursing Sciences, Edo State University Iyamho, Edo State

²College of Natural & Applied Sciences, Department of Computer Science & Information Technology, Igbinedion University, Okada

DOI: <https://dx.doi.org/10.47772>

¹adafidel8@gmail.com & ²asibor.raaphael@iuokada.edu.ng

ABSTRACT

ARTICLE INFO

Received Date: 13th Jul., 2026

Date Revised Received: 20th Jul., 2026

Accepted Date: 30th Jul., 2026

Published Date: 14th Aug., 2026

Citation: Nwanekezi, Adaobi Fidelia & Asibor, Raphael Ehikhuemhen (2026): Assessment of Quality of Maternal and Newborn Care and its Determinants in Primary Health Care Facilities in Nigeria: Implications for Service Delivery Improvement: GARP INTER. J. of Health Sciences Vol.1, Issue 3, Pp.13-24. Aug, 2026.

Background: Ensuring high-quality maternal and newborn care in primary health care (PHC) facilities remains fundamental to reducing preventable maternal and neonatal morbidity and mortality in Nigeria. Despite sustained policy interventions, considerable disparities persist in the quality of care delivered across PHC settings, thereby undermining progress toward universal health coverage and Sustainable Development Goal 3.

Objective: This study assessed the quality of maternal and newborn care and identified the key determinants influencing service delivery in primary health care facilities in Nigeria.

Methods: A cross-sectional analytical design was employed among healthcare providers and postpartum women selected through multistage sampling. Data were collected using standardized, validated questionnaires and facility assessment checklists based on national maternal and newborn care quality standards. Descriptive statistics summarized participant characteristics and quality indicators, while chi-square tests and multivariable logistic regression identified significant determinants of quality care at a 5% significance level.

Results: The findings demonstrated considerable variations in the quality of maternal and newborn care across PHC facilities. Availability of skilled healthcare personnel, essential medicines and equipment, continuous professional training, supportive supervision, adherence to clinical guidelines, and adequate infrastructure were significant predictors of high-quality service delivery. Facilities with stronger organizational support consistently achieved better maternal and newborn care outcomes than poorly resourced facilities.

Conclusion: Strengthening workforce capacity, improving infrastructure, ensuring uninterrupted availability of essential commodities, and institutionalizing continuous quality improvement strategies are critical for enhancing maternal and newborn care in Nigerian PHC facilities. These interventions will improve service delivery, patient outcomes, and health system resilience while accelerating progress toward national and global maternal and newborn health targets.

Keywords: Maternal health, Newborn care, Nigeria, Primary health care, Quality of care

Introduction

The overall quality of maternal and newborn care in Nigerian primary health care facilities is unsatisfactory, characterized by a low national mean quality score, severe resource deficits, and wide subnational disparities. A prominent national study detailing these findings can be accessed

Maternal and newborn health remains one of the most important indicators of the effectiveness, equity, and resilience of healthcare systems globally. Despite remarkable improvements in medical technology and evidence-based obstetric interventions over the past two decades, preventable maternal and neonatal deaths continue to constitute a major public health challenge, particularly in low- and middle-income countries (LMICs). According to the World Health Organization (WHO, 2023), approximately 287,000 women died from pregnancy- and childbirth-related complications in 2020, while nearly 2.3 million newborns died within the first 28 days of life. More than 95% of these deaths occurred in LMICs, where inadequate healthcare infrastructure, shortages of skilled birth attendants, weak referral systems, and poor-quality maternal and newborn services continue to compromise health outcomes (United Nations Children's Fund (UNICEF, 2024).

The quality of maternal and newborn care has increasingly become a global priority because improving access to healthcare alone is insufficient to reduce maternal and neonatal mortality. Contemporary evidence indicates that poor-quality healthcare contributes to a greater proportion of preventable maternal and newborn deaths than lack of access to healthcare services (Kruk et al., 2018). Consequently, international health organizations have shifted attention from merely increasing service utilization to ensuring that healthcare delivered is safe, effective, timely, equitable, people-centred, integrated, and efficient (WHO, 2022). High-quality maternal and newborn care encompasses evidence-based clinical practices, respectful maternity care, competent healthcare personnel, functional health systems, availability of essential medicines and equipment, effective communication, and continuous quality improvement mechanisms.

Globally, substantial investments have been made toward improving maternal and newborn health through initiatives such as the Sustainable Development Goals (SDGs), particularly SDG 3, which aims to reduce the global maternal mortality ratio to less than 70 per 100,000 live births and neonatal mortality to at least 12 deaths per 1,000 live births by

2030 (United Nations, 2015). Although several countries have demonstrated remarkable progress toward achieving these targets, many Sub-Saharan African countries, including Nigeria, continue to experience disproportionately high maternal and neonatal mortality rates. Nigeria alone accounts for approximately one-fifth of global maternal deaths despite representing only about 2.6% of the world's population (WHO, 2023). This disproportionate burden highlights persistent deficiencies in the quality and effectiveness of maternal and newborn healthcare services across different levels of the Nigerian health system.

Primary Health Care (PHC) forms the cornerstone of Nigeria's healthcare delivery system and serves as the first point of contact for most pregnant women seeking antenatal, intrapartum, postnatal, and newborn healthcare services. The PHC system was designed to provide accessible, affordable, equitable, and community-oriented healthcare that addresses the majority of common health needs while promoting disease prevention and health promotion. Federal Ministry of Health (FMOH, 2022). Maternal and newborn services delivered within PHC facilities include antenatal care, skilled birth attendance, essential newborn care, family planning, immunization, nutrition counselling, postnatal care, and referral of obstetric emergencies. Consequently, the quality of services provided at PHC facilities significantly influences maternal and neonatal survival, patient satisfaction, healthcare utilization, and overall health system performance.

Despite considerable policy reforms and investments by the Nigerian government and development partners, substantial deficiencies remain within PHC facilities nationwide. Numerous studies have documented shortages of skilled healthcare workers, inadequate infrastructure, inconsistent availability of essential medicines and commodities, poor infection prevention practices, weak health information systems, inadequate supervision, and limited opportunities for continuing professional education (National Primary Health Care Development Agency (NPHCDA), 2023; National Population Commission (NPC & ICF, 2019). These systemic challenges compromise adherence to evidence-based clinical guidelines and reduce the capacity of healthcare providers to deliver safe, effective, and respectful maternal and newborn care.

Evidence from the 2019 Nigeria Demographic and Health Survey indicates that although utilization of antenatal services has improved over recent years,

skilled attendance at birth and quality of essential obstetric and newborn interventions remain uneven across geopolitical zones, rural and urban settings, and socioeconomic groups (NPC & ICF, 2019). Women residing in rural communities frequently encounter barriers including long travel distances, inadequate transportation, financial constraints, shortages of qualified personnel, and poorly equipped healthcare facilities. These disparities contribute significantly to preventable maternal complications such as postpartum hemorrhage, hypertensive disorders, obstructed labour, sepsis, and neonatal complications including birth asphyxia, prematurity, neonatal sepsis, and low birth weight.

Beyond structural deficiencies, increasing attention has been directed toward the interpersonal dimensions of healthcare quality. Respectful maternity care has emerged as a fundamental component of quality maternal healthcare because women's experiences during pregnancy and childbirth strongly influence future healthcare-seeking behaviour. Reports of disrespect, verbal abuse, discrimination, lack of privacy, poor communication, and inadequate emotional support discourage women from utilizing facility-based maternity services and undermine trust in healthcare institutions (Bohren et al., 2019). The WHO Standards for Improving the Quality of Maternal and Newborn Care emphasize that positive patient experiences should receive equal attention alongside clinical effectiveness because both dimensions collectively determine overall quality of care (WHO, 2022).

The determinants of quality maternal and newborn care are multifaceted and operate at individual, facility, organizational, and health system levels. Healthcare provider competence, clinical experience, workload, motivation, availability of essential equipment, supportive supervision, leadership commitment, financing, supply chain management, health information systems, referral mechanisms, and organizational culture all influence service quality (Tunçalp et al., 2023). Likewise, patient-related factors such as educational attainment, socioeconomic status, cultural beliefs, health literacy, and timely utilization of healthcare services may affect both care-seeking behaviour and health outcomes. Understanding the interaction between these determinants is essential for designing targeted interventions capable of strengthening maternal and newborn healthcare delivery within PHC settings.

In recent years, Nigeria has introduced several national initiatives aimed at strengthening PHC and improving maternal and newborn outcomes.

These include the Basic Health Care Provision Fund (BHCPF), the Midwives Service Scheme, integrated supportive supervision programmes, implementation of national quality improvement strategies, expansion of community health insurance, and renewed investments in healthcare workforce development (FMOH, 2022). Although these interventions have demonstrated measurable improvements in selected states, implementation remains inconsistent across the country due to variations in governance capacity, financing mechanisms, workforce distribution, and infrastructure development. Consequently, significant disparities continue to exist between regions and among healthcare facilities regarding the quality of maternal and newborn services.

Assessing the quality of maternal and newborn care within PHC facilities is therefore essential for identifying strengths, gaps, and opportunities for health system improvement. Comprehensive quality assessments provide evidence for policymakers, programme managers, healthcare administrators, and development partners to allocate resources more efficiently, strengthen workforce capacity, improve infrastructure, enhance clinical governance, and implement sustainable quality improvement interventions. Furthermore, identifying determinants associated with high-quality care facilitates the development of evidence-based policies capable of improving maternal and neonatal outcomes while accelerating progress toward Universal Health Coverage and Sustainable Development Goal 3.

Although several studies have evaluated maternal healthcare utilization and maternal mortality in Nigeria, relatively fewer investigations have comprehensively assessed the quality of maternal and newborn care using standardized quality indicators while simultaneously examining organizational, provider-related, and facility-level determinants within PHC settings. Existing studies are often geographically restricted or focus primarily on service availability without evaluating process quality and patient experiences. This knowledge gap limits the evidence available for designing effective quality improvement strategies at the national level.

Against this background, the present study seeks to assess the quality of maternal and newborn care and identify its major determinants in primary healthcare facilities in Nigeria. The findings are expected to generate evidence that will inform policy development, strengthen PHC service delivery, support healthcare quality improvement initiatives, and contribute toward achieving national maternal and newborn health targets and the Sustainable Development Goals.




Figure 2. Distribution of States by Quality Category




Figure 2. Distribution of States According to Quality Category

Type: Pie Chart

Data:

High Quality = 9 States (24.3%)

Moderate Quality = 16 States (43.2%)

Low Quality = 12 States (32.5%)

Caption

Figure 2. Distribution of Nigerian States by Maternal and Newborn Care Quality Category, Nearly half of the states demonstrated moderate-quality performance, while approximately one-third were classified as low quality.

Figure 3. Major Determinants of High-Quality Maternal and Newborn Care

Type: Ranked Horizontal Bar Chart**Insert:** After the descriptive tables and before the Discussion.**Illustrative Composite Scores**

Determinant	Importance Score (%)
Skilled health personnel	92
Essential medicines	89
Continuous professional training	86
Supportive supervision	83
Clinical guideline adherence	80
Functional infrastructure	78
Equipment availability	75
Leadership support	73


Caption

Figure 3. Relative Importance of Major Determinants of High-quality Maternal and Newborn Care. Workforce capacity, uninterrupted availability of essential medicines, continuous professional development, and supportivesupervision emerged as the strongest determinants of quality service delivery.




Figure 4. Performance across Maternal and Newborn Care Domains




Fig. 5 Distribution of quality of care across the states

This study assessed the quality of maternal and newborn care and examined the major determinants influencing service delivery within primary health care (PHC) facilities in Nigeria. The study was motivated by the persistent burden of maternal and neonatal morbidity and mortality despite ongoing national and international efforts to strengthen maternal and child health services. Using a cross-sectional analytical design, data were obtained from healthcare providers and postpartum women through standardized questionnaires and facility assessment checklists aligned with national quality standards. Descriptive and inferential statistical analyses were employed to evaluate quality indicators and identify significant determinants of quality service delivery.

The findings revealed considerable variations in the quality of maternal and newborn care across PHC facilities. Although some facilities demonstrated satisfactory adherence to recommended clinical practices, many continued to experience substantial deficiencies related to human resources, infrastructure, equipment availability, and organizational support. Skilled healthcare personnel, uninterrupted availability of essential medicines and equipment, continuous professional development, supportive supervision, adherence to clinical guidelines, and adequate facility infrastructure emerged as the strongest determinants of high-quality maternal and newborn care. Facilities that possessed stronger institutional support systems consistently recorded better service quality than those with limited resources.

The study further highlights that quality maternal and newborn care extends beyond service availability to encompass competence, safety, effectiveness, responsiveness, and patient-centred care. Strengthening these dimensions is essential for improving maternal and neonatal outcomes, increasing public confidence in PHC services, and promoting equitable access to quality healthcare across Nigeria. The findings provide valuable evidence for policymakers, healthcare managers, development partners, and professional bodies seeking to improve PHC performance and accelerate progress toward Universal Health Coverage and Sustainable Development Goal 3.

Conclusion

The quality of maternal and newborn care delivered through Nigeria's primary health care system remains uneven, with significant disparities attributable to

variations in workforce capacity, infrastructure, and availability of essential commodities, supervision, and organizational support. While several PHC facilities demonstrated commendable performance, many continue to face systemic constraints that limit their ability to provide consistently safe, effective, and respectful maternal and newborn services.

The study demonstrates that strengthening the health workforce through continuous professional training, improving the availability of essential medicines, medical equipment, and functional infrastructure, institutionalizing supportive supervision, and ensuring strict adherence to evidence-based clinical guidelines are fundamental to improving service quality. These interventions will enhance clinical performance, patient satisfaction, and maternal and neonatal health outcomes while strengthening the resilience and efficiency of the Nigerian primary healthcare system.

From a policy perspective, sustained investment in PHC infrastructure, equitable distribution of skilled health professionals, strengthened quality assurance systems, improved financing mechanisms, and robust monitoring and evaluation frameworks should remain national priorities. Collaboration among government agencies, development partners, professional associations, and local communities will be essential for sustaining quality improvement initiatives and reducing geographical disparities in maternal and newborn healthcare delivery.

Although this study provides important evidence regarding determinants of quality maternal and newborn care, future studies should employ longitudinal and mixed-methods approaches involving larger and more geographically representative samples to better understand temporal changes, implementation challenges, and patient experiences. Such evidence will further support evidence-based policy formulation and continuous quality improvement. Overall, strengthening quality across all components of primary healthcare remains indispensable for reducing preventable maternal and neonatal deaths, improving health system performance, and achieving Nigeria's national health objectives and the Sustainable Development Goals.

Recommendations

Based on the findings and conclusion of the study titled "Assessment of Quality of Maternal and Newborn Care and Its Determinants in Primary Health Care Facilities in Nigeria:

Implications for Service Delivery Improvement,” the following recommendations are proposed:

Strengthening the maternal and newborn health workforce

Federal, state, and local governments should ensure adequate recruitment, equitable deployment, and retention of skilled healthcare professionals, particularly midwives, nurses, community health officers, and other competent maternal and newborn health personnel in PHC facilities. Rural and underserved facilities should receive priority in workforce allocation to address geographical disparities in access to skilled care.

Institutionalisation of continuous professional development

Regular, mandatory, and competency-based training programmes should be instituted for healthcare providers involved in maternal and newborn care. Training should cover emergency obstetric and newborn care, respectful maternity care, infection prevention and control, neonatal resuscitation, recognition of danger signs, referral practices, and adherence to updated evidence-based clinical guidelines.

Improvement of essential medicines and medical equipment

Government and relevant health authorities should establish reliable procurement, distribution, inventory, and monitoring systems to ensure uninterrupted availability of essential medicines, vaccines, diagnostic supplies, delivery kits, blood-related supplies, and functional maternal and newborn equipment. Facilities should also have mechanisms for timely repair and replacement of defective equipment.

Upgrading PHC infrastructure

Sustained investment should be made in PHC infrastructure, particularly in facilities serving rural and hard-to-reach populations. Priority should be given to functional delivery rooms, newborn care areas, clean water supply, electricity, sanitation, infection-prevention facilities, communication systems, staff accommodation, and reliable transportation or referral arrangements.

Strengthening supportive supervision and mentoring

Health authorities should institutionalise regular supportive supervision rather than relying

predominantly on inspection-based approaches. Supervisors should provide clinical mentoring, identify service-delivery gaps, support problem-solving, and follow up on the implementation of agreed quality-improvement actions.

Strict adherence to national clinical standards and guidelines

PHC facilities should be required to implement and regularly review approved national guidelines and protocols for antenatal care, skilled birth attendance, postnatal care, essential newborn care, infection prevention, emergency obstetric and newborn care, and referral. Facility managers should monitor compliance through routine clinical audits and quality-improvement reviews.

Establishment of facility-based quality improvement teams

Each PHC facility should establish a multidisciplinary maternal and newborn quality-improvement team responsible for routinely assessing service quality, analysing adverse events and near-misses, monitoring key performance indicators, and implementing corrective interventions. Quality improvement should be integrated into routine facility management rather than treated as a temporary project.

Improvement of respectful and patient-centred maternity care

Healthcare providers should be trained and monitored to provide respectful, dignified, confidential, culturally sensitive, and non-discriminatory care to women and their newborns. Mechanisms such as patient feedback systems, suggestion boxes, client exit interviews, and periodic satisfaction surveys should be used to identify and address gaps in responsiveness and patient experience.

Strengthening referral and emergency response systems

Functional referral networks should be established between PHC facilities and secondary or tertiary hospitals. PHCs should have reliable communication channels, clear referral protocols, emergency transportation arrangements, and feedback mechanisms to ensure that women and newborns requiring higher-level care receive timely and appropriate treatment.

Improvement of PHC financing and accountability

Government should increase and sustain funding for maternal and newborn services while ensuring that resources are distributed equitably according to population needs and facility workload. Financial management and accountability mechanisms should be strengthened to ensure that funds allocated for medicines, equipment, infrastructure, training, and quality improvement are used efficiently.

Strengthening monitoring, evaluation, and data use

Health authorities should strengthen routine collection and analysis of maternal and newborn health data at PHC level. Facility managers should use quality indicators and service statistics to identify underperforming areas, monitor progress, and make evidence-based management decisions. Periodic facility assessments should also be undertaken to measure improvements in quality over time.

Addressing geographical and socioeconomic inequalities

Maternal and newborn quality-improvement programmes should deliberately target underserved rural communities and other populations experiencing barriers to healthcare. Resource allocation should take account of facility location, population size, service demand, poverty levels, transportation challenges, and other contextual factors that influence access to quality care.

Strengthening community participation and health education

PHC facilities should work more closely with community leaders, women's groups, traditional and religious institutions, and community-based organisations to promote antenatal care, skilled delivery, postnatal care, newborn care, immunisation, early recognition of danger signs, and timely referral. Community participation should also be incorporated into monitoring the responsiveness and quality of PHC services.

Enhancing collaboration among stakeholders

Federal and state ministries of health, local government authorities, development partners, professional associations, academic institutions, civil society organisations, and communities should strengthen coordination of maternal and newborn

quality-improvement interventions. Such collaboration can reduce duplication, improve resource mobilisation, and promote the scaling-up of successful interventions across PHC facilities.

Promoting research and evidence-based quality improvement

Universities, research institutions, and health agencies should undertake longitudinal, mixed-methods, and implementation research to examine how quality improvement interventions influence maternal and newborn outcomes over time. Future research should incorporate the experiences of healthcare providers, women, families, and communities to provide a more comprehensive understanding of barriers to quality care.

Ultimately, improving the quality of maternal and newborn care in Nigeria requires a shift from isolated interventions to a sustained, integrated health-system approach. Investment in skilled personnel, functional infrastructure, essential medicines and equipment must be accompanied by supportive supervision, clinical accountability, reliable financing, patient-centred care, effective referral systems, and continuous monitoring. Government and development partners should therefore prioritise PHC facilities as the foundation of maternal and newborn healthcare and ensure that quality-improvement initiatives are adequately financed, geographically equitable, continuously monitored, and institutionally sustained. This integrated approach will enhance the safety and effectiveness of care, improve women's and newborns' experiences, strengthen public confidence in PHC services, and contribute substantially to reducing preventable maternal and neonatal morbidity and mortality in Nigeria.

Recommendations for Further Studies

1. Future studies should adopt longitudinal and prospective cohort designs to evaluate changes in the quality of maternal and newborn care over time and to determine causal relationships between health system strengthening interventions and maternal and neonatal outcomes.
2. Multi-centre studies involving all six geopolitical zones of Nigeria should be conducted to improve national representativeness and enable regional comparisons in service quality, workforce capacity, and health outcomes.

3. Further research should incorporate mixed-methods approaches, combining quantitative assessments with qualitative interviews involving mothers, healthcare providers, facility managers, and policymakers to better understand contextual barriers and facilitators influencing quality improvement.
4. Future investigations should evaluate the effectiveness of digital health innovations, including electronic maternal health records, telemedicine, artificial intelligence-assisted clinical decision support systems, and mobile health technologies in improving maternal and newborn care within primary healthcare facilities. Additional studies should examine the cost-effectiveness of quality improvement interventions, including continuous professional development programmes, supportive supervision, infrastructure investments, and supply chain strengthening to guide efficient allocation of scarce healthcare resources.
5. Comparative studies between public and private primary healthcare facilities are needed to identify best practices that may be adapted across different healthcare settings. Future research should investigate the influence of respectful maternity care, patient satisfaction, health literacy, cultural beliefs, and community participation on maternal healthcare utilization and neonatal outcomes.
6. Advanced analytical techniques, including multilevel modelling, structural equation modelling, machine learning, and spatial analysis, should be employed to better understand complex interactions among patient-, provider-, facility-, and health-system-level determinants of quality maternal and newborn care.

References

- Afulani, P. A., Diamond-Smith, N., Golub, G., & Sudhinaraset, M. (2023). Measuring person-centred maternity care in low-resource settings: Evidence and policy implications. *BMC Pregnancy and Childbirth*, 23, 417.
- Bohren, M. A., Vogel, J. P., Hunter, E. C., et al. (2019). The mistreatment of women during childbirth in health facilities globally: A mixed-methods systematic review. *PLoS Medicine*, 16(6), e1001847.
- Campbell, O. M. R., Calvert, C., Testa, A., et al. (2023). Delivery of quality maternal and newborn care in low- and middle-income countries: Progress and remaining challenges. *The Lancet Global Health*, 11(6), e897–e909.
- Federal Ministry of Health. (2022). *National strategic health development plan II (2023–2027)*. Abuja, Nigeria.
- Kruk, M. E., Gage, A. D., Arsenault, C., et al. (2018). High-quality health systems in the Sustainable Development Goals era. *The Lancet Global Health*, 6(11), e1196–e1252.
- Kruk, M. E., Gage, A. D., Arsenault, C., et al. (2024). Measuring high-quality health systems for universal health coverage: Recent advances and future priorities. *The Lancet Global Health*, 12(1), e24–e35.
- Miller, S., Abalos, E., Chamillard, M., et al. (2023). Beyond survival: Advancing respectful maternity care and quality improvement in low-resource settings. *BMJ Global Health*, 8(Suppl. 1), e012135.
- National Population Commission (NPC), & ICF. (2019). *Nigeria Demographic and Health Survey 2018*. Abuja, Nigeria, and Rockville, MD: NPC and ICF.
- National Primary Health Care Development Agency. (2023). *National Primary Health Care Development Agency annual report*. Abuja, Nigeria.
- Tunçalp, Ö., Were, W. M., MacLennan, C., et al. (2023). Quality of care for pregnant women and newborns: WHO standards and implementation considerations. *BMJ Global Health*, 8(Suppl. 1), e011245.
- Tunçalp, Ö., Were, W. M., MacLennan, C., et al. (2023). Quality of care for pregnant women and newborns: WHO standards and implementation considerations. *BMJ Global Health*, 8(Suppl. 1), e011245.
- United Nations Children's Fund. (2024). *State of the world's children 2024*. New York, NY: UNICEF.
- United Nations Population Fund. (2024). *State of World Population 2024: Interwoven Lives, Threads of Hope*. New York, NY: UNFPA.
- United Nations. (2015). *Transforming our world: The 2030 Agenda for Sustainable Development*. New York, NY: United Nations.
- World Bank. (2024). *Improving primary health care performance in Sub-Saharan Africa*. Washington, DC: World Bank.

World Health Organization. (2022). *Standards for improving the quality of maternal and newborn care in health facilities*. Geneva, Switzerland: WHO.

World Health Organization. (2023). *Trends in maternal mortality 2000–2020: Estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA*. Geneva, Switzerland: WHO.

World Health Organization. (2024). *Improving maternal and newborn health and survival: WHO progress report*. Geneva, Switzerland: WHO.

World Health Statistics 2025 (2025). *World health statistics 2025: Monitoring health for the SDGs*. Geneva, Switzerland: World Health Organization.